Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023



## **NHS LOTHIAN**

# DIRECTORATE OF LABORATORY MEDICINE

# Scottish Mycobacteria Reference Laboratory

# & NHS Lothian Mycobacteria Laboratory

# **User Manual**

Authors: Dr I.F. Laurenson & Benjamin Moore

Scottish Microbiology Reference Laboratories, Edinburgh:

http://www.edinburghlabmed.co.uk/Specialities/reflab/Pages/default.aspx

Authority For Issue: Benjamin Moore	Page 1 of 24
Document printed from Q-pulse 16/1	0/2023 09:28:00 by benjamin.moore

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

### UKAS ISO 15189:2012 Accredited

#### INDEX

Scottish Mycobacteria Reference Laboratory, Scottish Reference Laboratories, Edinburgh	2
Telephone and Email Enquiries	3
Core Hours	3
Contact List (Permanent Staff)	4
Packing and Transport of Infectious Substances	5
Levels of Service	7
Samples for Mycobacterial Investigations	8
Sample Types and Containers for Mycobacteria Investigation	12
Tests Available, Turn Around Times and Costs	14
Molecular Detection of Mycobacteria inClinical Specimens	17
Reporting of Results	18
Technical, Scientific and Health and Safety Advice	21
Clinical, Epidemiological and Infection Control Advice	21
Training and Education	21
Research and Audit	22
Quality Assurance	22
Protection of Personal Information	22
Antimicrobial Assays	23
Leprosy Diagnosis	24

Authority For Issue: Benjamin Moore	Page 2 of 24
Document printed from Q-pulse 16/1	0/2023 09:28:00 by benjamin.moore

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

#### Scottish Mycobacteria Reference Laboratory, Scottish Reference Laboratories, Edinburgh

The Scottish Mycobacteria Reference Laboratory (SMRL) is the Scottish Reference Laboratory for Mycobacteriology, particularly for the identification, drug susceptibility testing and strain typing of *Mycobacterium tuberculosis* complex (MTBC) and non-tuberculous mycobacteria (NTM). The laboratory receives approximately 1600 mycobacterial cultures per year mainly from Scottish NHS laboratories. The service is also funded to perform rapid detection of mycobacteria and screening for drug resistance on acid-fast bacillus (AFB) smear positive pulmonary samples although this function is expected to be superseded, by each Scottish Health Board laboratory. Further information regarding advice available is detailed below. SMRL is also the Scottish repository for mycobacterial cultures.

The laboratory is based within the Directorate of Laboratory Medicine, Royal Infirmary of Edinburgh, Little France Crescent, Edinburgh, EH16 4SA, UK. It is part of the Scottish Microbiology Reference Laboratories, Edinburgh and is fully integrated with the combined laboratories of NHS Lothian. SMRL is funded through an agreement between National Service Scotland and NHS Lothian. Some services have been developed which are available at a charge to the user.

This manual covers three levels of service:

- 1. National Reference services funded by National Services Scotland (NSS).
- 2. Services paid for by charging the users per test.
- 3. Services provided by NHS Lothian to the local and other Boards by Service Level Agreement. These Boards are currently, NHS Borders, NHS Fife, NHS Dumfries and Galloway, NHS Highland, NHS Forth Valley and NHS Tayside.

#### **Telephone and Email Enquiries**

TB Laboratory Tel: 0131 242 6022/23 TB Office Tel: 0131 242 6016 The preferred method of communication for non urgent matters is Loth.Smrl@nhslothian.scot.nhs.uk

Fax machines are not considered a safe method of transmitting sensitive information; therefore, fax provision in NHS Lothian is not available.

#### **Core Hours**

The core laboratory hours are 9am to 5pm, Monday to Friday, but not weekends, Christmas and New Year holidays. At other times, Lothian laboratory users can discuss urgent AFB microscopy via the Consultant/Specialist Registrar on-call for clinical bacteriology (contact Royal Infirmary of Edinburgh switchboard Tel: 0131 536 1000). There is currently no out-of-hours AFB screening or Reference Laboratory service.

Authority For Issue: Benjamin Moore	Page 3 of 24
Document printed from Q-pulse 16/1	0/2023 09:28:00 by benjamin.moore
Document printed from Q-pulse 10/1	0/2023 03.20.00 by benjamin.moore

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

For urgent enquiries out-of-hours, the on call Consultant Microbiologist may be contacted through the RIE switchboard (Tel 0131 536 1000) in the first instance.

### **Contact List (Permanent Staff)**

Name and Designation	Telephone Numbers and e-mail Address
For non urgent enquiries	Loth.Smrl@nhslothian.scot.nhs.uk
	Tel 0131 242 6016
Mrs Mori Robertson, SMRL Administrator	Mori.Robertson@nhslothian.scot.nhs.uk
	Tel: 0131 242 6016
Dr Ian Laurenson, Director	lan.Laurenson@nhslothian.scot.nhs.uk
	Tel: 0131 242 6079
Dr Olga Moncayo, Deputy Director	Olga.Moncayo@nhslothian.scot.nhs.uk
	Tel: 0131 242 6070
Dr Donald Inverarity, Deputy Director	Donald.Inverarity@nhslothian.scot.nhs.uk
	Tel: 0131 242 6082
Dr Louise Seagar, Clinical Scientist	Louise.Seagar@nhslothian.scot.nhs.uk
	Tel: 0131 242 6009
Dr Naomi Gadsby, Clinical Scientist	Naomi.Gadsby@nhslothian.scot.nhs.uk
	Tel: 0131 242 1260
Mr Ben Moore, Clinical Scientist	Benjamin.Moore@nhslothian.scot.nhs.uk
	Tel: 0131 242 6009
Mrs Pauline Claxton, BMS Team Manager	Pauline.Claxton@nhslothian.scot.nhs.uk
	Tel: 0131 242 6009
Ms Ania Pilarska, BMS Team Manager	Anna.Pilarska@nhslothian.scot.nhs.uk
	Tel: 0131 242 6009
Ms Iryna Andrade, BMS Team Manager	Iryna.Andrade@nnsiotnian.scot.nns.uk
	Tel: 0131 242 6009
BMS Specialists:	
Ms Sandra Houston	
Mr Jack Oliver	
Mr Noil McMipp, MTO	
Mrs Rachel Rudge	
Mr Darren Robertson	Tal: 0121 242 6022 (6022
	181. 0131 242 0022/0023

Authority For Issue: Benjamin Moore	Page 4 of 24
Document printed from Q-pulse 16/10/2023 09:28:00 by benjamin.moore	
This is a controlled document. This copy is valid on day of print only after which the user must oncure that this is the c	

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

# The service is managed overall by NHS Lothian. If we are unable to deal with an issue and you wish to discuss a matter with NHS Lothian Management then contact:

Linda Mulhern Tel: 0131 242 6017 <u>linda.mulhern@nhslothian.scot.nhs.uk</u> Operational Science Manager, Microbiology, NHS Lothian

Amanda Malham Tel: 0131 242 7051 <u>Amanda.Malham@nhslothian.scot.nhs.uk</u> Healthcare Science Manager, Cell Sciences, NHS Lothian.

#### **Packing and Transport of Infectious Substances**

#### **Postal Address**

Scottish Mycobacteria Reference Laboratory Scottish Microbiology Reference Laboratories, Edinburgh, Department of Laboratory Medicine Royal Infirmary of Edinburgh 51 Little France Crescent Old Dalkeith Road Edinburgh EH16 4SA DX Address

Exchange: Edinburgh 96 EH DX Number: DX6231201

NB Infectious substances may only be transported in packaging, which meets the U.N. Class 6.2 specifications and the 602 packing requirements. All Cultures must be sent by DX Courier or equivalent-NOT by post. They must conform to the Transport of Dangerous Goods Act. Cultures for 'routine' processing should not usually be sent in batches.

Packaging Mycobacteria Cultures or Samples:

- 1. Remove any water of condensation from the solid culture vial(s) taking normal safety precautions for handling cultures.
- 2. Seal the cap with 'parafilm'.
- 3. Wrap individual vials in sufficient absorbent material to absorb all the liquid in the vial and to protect it from breakage.
- 4. Seal this package twice in plastic bags.

#### For posting by DX courier for all cultures – preferred for specimens:

Authority For Issue: Benjamin Moore	Page 5 of 24
Document printed from Q-pulse 16/1	0/2023 09:28:00 by benjamin.moore
This is a controlled document: This copy is valid on day of	of print only, after which the user must ensure that this is the cor

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

- 1. Place the sealed specimen bag into a screw-cap container from a DX box and put with it the corresponding request forms.
- 2. Carefully place the destination address label provided on the green tracking label on the package, taking care not to cover any of the peel-off barcodes or tracking numbers. Also leave some space next to the lines for receiver's DX number and Exchange code. Place some clear sellotape over the label to ensure it is not easily removed or damaged by rain, again avoiding covering any peel-off labels.
- 3. Fill in the DX number and Exchange code of the destination.
- 4. Peel off <u>one</u> of the two matching tracking numbers at the bottom of the Tracked Specimen label and stick it in the provided DX logbook and fill in the receiver's details and date of dispatch.
- 5. At the top of the DX label, fill in the sender's name, sender's DX number, sender's exchange code and the name and contact number for the sender. This information is written on a separate card at the front of the DX logbook.
- 6. Place the provided blue security seal sticker over the package closure and dispatch.

#### For posting by First Class Post (Specimens Only):

- 1. Place the wrapped sample inside the inner container of the packaging and tighten the lid.
- 2. Place the inner container into the outer cardboard box.
- 3. Close the box and secure with tape.
- 4. Stick address label on the top of the box.
- 5. Stick "From" label, with senders address, on the bottom of the box.
- 6. Post by <u>First Class Post</u> or courier such as DX for samples.

Authority For Issue: Benjamin Moore	Page 6 of 24
Document printed from Q-pulse 16/1	0/2023 09:28:00 by benjamin.moore

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

#### **Levels of Service**

#### **1.** National Services Scotland funded Remit

SMRL receives cultures of presumed mycobacteria from all NHS laboratories in Scotland. These cultures are tested to determine

- Presence of mycobacteria.
- Full species identification.
- Susceptibility to standard anti-mycobacterial drugs.
- Susceptibility to second line anti-mycobacterial drugs
- Genomic comparison of *M. tuberculosis* complex strains.
- Use of molecular tests on smear positive respiratory samples to rapidly identify the presence of mycobacteria and screen for rifampicin drug resistance in *M.tuberculosis* complex.

#### 2. Services available by charging users.

- Rapid liquid culture of samples from normally sterile sites (e.g. blood, CSF, bone marrow, tissue samples, aspirates and bronchoalveolar lavage samples).
- Molecular tests on smear negative samples and smear positive non-respiratory samples.
- Molecular tests to screen for isoniazid, fluoroquinolones, amikacin, capreomycin, kanamycin and ethionamide drug resistance in isolates of *M. tuberculosis* complex.
- Culture of smear negative samples.
- Identification of slow growing mycobacterial species from environmental water and air samples e.g. from heater-cooler units and environments where these units are in operation.

#### 3. Services available by contract between requesting service and NHS Lothian

• Processing of routine samples including blood cultures from patients for mycobacteria investigation.

Please note SMRL no longer offers PCR for MTBC detection in formalin-fixed, paraffin embedded tissue. Samples for 16S rDNA PCR may be sent directly by the user to a suitably accredited laboratory such as:

Microbiology, Virology and Infection Control Level 4, Camelia Botnar Laboratories Great Ormond Street Hospital for Children Great Ormond Street London WC1N 3JH DX number: DX6640203 DX Exchange: Bloomsbury 91WC http://www.labs.gosh.nhs.uk/laboratory-services/microbiology-virology-and-infection-

#### control/tests/16s-rdna-pcr

For histological samples requests a minimum of 4 rolled sections (10µm thick). More preferred if possible to allow repeat extractions.

Authority For Issue: Benjamin Moore	Page 7 of 24
Document printed from Q-pulse 16/1	0/2023 09:28:00 by benjamin.moore
This is a controlled decument: This copy is yelid on day a	f print only often which the year must ensure that this is the

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

Department of Microbiology Leeds General Infirmary Old Medical School Thoresby Place Great George Street Leeds LS1 3EX DX number: DX6281504 DX Exchange: Leeds 90LS https://www.leedsth.nhs.uk/a-z-of-services/pathology/test-and-tubes/molecularbacteriology/mycobacterial-non-tb-and-tb-pcr For histological samples requests 10 sections 10µm thick

#### Samples for Mycobacterial Investigations

#### 1. Microscopy

All samples except blood, swabs and urine are routinely examined for direct AFB microscopy (provided there is sufficient specimen).

#### 2. Culture

Our current practice is to inoculate all specimens into liquid and solid media.

#### 3. Drug susceptibility testing

Susceptibilities are routinely performed on the first MTB complex isolate from each patient and then only repeated at 3 month intervals.

NTM susceptibility testing is based on the ATS/IDSA criteria for diagnosing NTM infection, modified by local experience and thereafter only repeated at 3 month intervals. Testing will be performed on:

- all NTM from sterile sites (except for *Mycobacterium marinum* in skin);
- first isolates of *Mycobacterium abscessus* complex, *Mycobacterium xenopi, Mycobacterium malmoense and Mycobacterium kansasii*. Experience suggests that these are likely to be associated with clinically significant disease
- patients with cystic fibrosis, HIV or with significant T cell immunosuppression (if we are informed of the clinical features)

For all other pulmonary isolates of NTM we will not do sensitivity testing unless we are informed that the patient meets ATS criteria for NTM lung disease (please indicate which criteria are met). For non sterile sites such as urine or faeces we will not do sensitivity testing unless there is a strong clinical suspicion of infection. To arrange sensitivity testing please email us at: Loth.Smrl@nhslothian.scot.nhs.uk

Authority For Issue: Benjamin Moore	Page 8 of 24
Document printed from Q-pulse 16/1	0/2023 09:28:00 by benjamin.moore

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

The current BTS NTM Pulmonary Disease advice, including comment on ATS/IDSA criteria is at: <u>https://www.brit-thoracic.org.uk/standards-of-care/guidelines/bts-guidelines-for-non-tuberculous-mycobacteria</u> (accessed January 2022).

#### 4. Molecular tests

Pulmonary specimens from NHS Lothian sources fulfilling NICE based criteria will be tested by a rapid molecular test for MTB complex and rifampicin resistance if sufficient sample is received (2ml minimum volume required). In addition, any AFB smear positive samples are also tested. Rapid molecular testing of both pulmonary specimens and positive cultures is also available for MTB complex and isoniazid, fluoroquinolone, amikacin, kanamycin, capreomycin and ethionamide resistance where clinically indicated.

#### 5. Whole Genome Sequencing (WGS) of mycobacterial cultures

The SMRL is currently using WGS for all mycobacterial species identifications, resistance prediction of MTB complex and genetic relatedness of *M.tuberculosis* complex isolates from positive cultures. Rapid confirmation of MTB complex is achieved initially using lateral-flow immunochromatographic assay. Phenotypic susceptibility testing may be used to confirm MTB complex susceptibility pattern results if drug resistance is suspected.

There may occasionally be situations where cultures are referred for alternative modes of testing at one of the PHE National Mycobacterium Reference Service laboratories.

Users should inform us in advance of cultures that are being sent by post or courier preferably by 'generic' e-mail: <u>Loth.Smrl@nhslothian.scot.nhs.uk</u>. We will issue a receipt report within one working day from receipt of cultures.

For urgent requests or for direct molecular detection add-on testing, a SMRL staff member should be contacted by phone (see contact list above).

A completed SMRL request form **must** accompany every culture or specimen. <u>http://www.edinburghlabmed.co.uk/Specialities/reflab/Pages/default.aspx</u>

Wherever possible, please time the posting of samples and cultures to minimise the possibility of these lying unattended over weekends and holidays. The SMRL will attempt to process all primary samples and referred cultures as much as possible. All required information on the sample or request form that is missing could result in a delay in sample processing.

All requests must provide the following Minimum Data Set information:

#### Forms

- 1. Patient Identifier Number (CHI-for Scottish patients)
- 2. Postcode of patient where available
- 3. Sender's Laboratory Number
- 4. Surname
- 5. Forename

Authority For Issue: Benjamin Moore	Page 9 of 24
Description of the second seco	
Document printed from Q-pulse 16/1	0/2023 09:28:00 by benjamin.moore
This is a controlled document: This copy is valid on day of	of print only, after which the user must ensure that this is the

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

- 6. Date of Birth
- 7. Gender
- 8. Location of Sender
- 9. Specimen type (or, if culture isolate, specimen type from which the culture was isolated).
- 10. Any relevant clinical information. If results need to be telephoned please supply contact details.
- 11. AFB and molecular results were performed at sending laboratory
- 12. If possible patient's hospital and consultant.

#### Sample

Minimum data are also required on the sample container:

- 1. Surname
- 2. Forename
- 3. Date of Birth
- 4. Sample date and time
- 5. Sample type and site
- 6. Location of patient
- 7. Sending Laboratory number if applicable

#### Vials

On culture vials, patient name and the sender's Laboratory Number are required as a minimum.

Specimens should be correctly collected and delivered as quickly as possible to the laboratory. Although *Mycobacteria tuberculosis* may survive in sputum for one week, the probability of successfully culturing AFB bacilli decreases with time and is especially critical for paucibacillary specimens. If specimens cannot be transported to the laboratory within one hour, it is recommended to store them at 4 °C. This does not apply to whole blood or bone marrow specimens in Myco/F Lytic bottles which should not be refrigerated or incubated prior to sending. Leaking cultures will <u>not</u> be processed.

#### 6. Interferon Gamma Release Assay (IGRA) for detection of latent TB infection.

Testing is currently performed 'in-house' for NHS Lothian patients by Blood Sciences Autocore using the QuantiFERON-TB Gold Plus (QFT-Plus) test. Other Health Boards may use this facility via a Service Level Agreement with NHS Lothian. Potential users should contact:

Mrs Una Jarrold Tel: 0131 242 6830 <u>Una.White@nhslothian.scot.nhs.uk</u>

Operational Science Manager, Blood Sciences, NHS Lothian.

Quantiferon 4 tube system and Sarstedt universal blood culture adaptors are available from catalogue on PECOS system for NHS Lothian users. If a Safety-Multifly is used, the line must be primed with an empty Sarstedt tube first to ensure the correct volume (1ml) of blood is delivered into each tube. All 4 tubes (Nil, TB1, TB2, and Mitogen) must be inoculated. The black mark on the

Authority For Issue: Benjamin Moore	Page 10 of 24
Document printed from Q-pulse 16/7	10/2023 09:28:00 by benjamin.moore

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

side of the tubes indicates the validated range of 0.8–1.2 ml. If the level of blood in any tube is outside the range of the indicator mark, a new blood sample should be obtained. It is important to gently agitate the tubes ten times to ensure the entire inner surface of each tube is coated with blood, to dissolve antigens on tube walls. DO NOT shake vigorously.

Samples should be dispatched to the laboratory as soon as possible but must reach the laboratory within 16 hours of collection. Do not refrigerate. See the Guidelines for NHS Lothian users of Microbiological and Virological Specialist laboratory for full details. Laboratories out with NHS Lothian may incubate tubes overnight at 37°C which may then be sent according to SLA to the NHS Lothian Specialist Serology/Virology centre.

The T-spot assay is another IGRA, which is available from:

Oxford Immunotec 115D Milton Park Abingdon Oxfordshire OX14 4RZ Tel: +44 (0) 1235 442 780 Fax: +44 (0) 1235 442 781 www.oxfordimmunotec.com

Note that tuberculin skin testing is an alternative mode of assessing latent TB infection (LTBI). There is no gold standard for LTBI detection. All tests may be negative in patients with active TB infection and LTBI tests have little role in routine assessment of active TB infection.

Authority For Issue: Benjamin Moore		Page 11 of 24
Document printed from Q-pulse 16/1	0/2023 09:28:00 by benjamin.moore	

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

## Sample Types and Containers for Mycobacteria Investigation

Specimen type	Containers and transport	Comments
Sputum (3 to preferably 5ml).	Disposable plastic, sterile "universal" container. Collected approximately 8-24 hours apart. Early morning samples preferred before ingestion of food	Decontaminated and neutralised samples may lose viability during transit and should not be sent
Induced sputum	Disposable plastic, sterile "universal" container	
Bronchoalveolar lavage (BAL)/ Bronchial Washings (3- 25ml)	Disposable plastic, sterile "universal" container	
Pleural fluids & aspirates (up to 25ml)	Disposable plastic, sterile "universal" container	Pleural and pericardial fluids are insensitive samples. Concurrent pleural or pericardial biopsy taken with the fluid is more useful
Biopsy material (including pleural biopsies)	Disposable plastic, sterile "universal" container (add a few drops of preservative free 0.9% sterile saline to keep sample moist)	DO NOT USE FORMOL SALINE OR FORMALIN
Tissues (including heart valves)	Disposable plastic, sterile "universal" container (add a few drops of preservative free 0.9% sterile saline to keep sample moist)	DO NOT USE FORMOL SALINE OR FORMALIN
Early Morning Urine	Sterile 500ml bottles filled up to 2/3 full. Send three early morning samples on successive days. Do not use bottles with boric acid. Midstream EMU is an acceptable but not ideal alternative.	EMU bottles available on request from the stores department, Astley Ainsley Hospital, Edinburgh for Lothian users.
Blood (1-5ml, optimum 3-5ml))	Myco/ F lytic blood culture bottles available from SMRL. Citrated blood also acceptable alternative at present.	Do not refrigerate or pre- incubate inoculated bottles.
Bone marrow	Use Myco/ F lytic blood culture bottles available from SMRL, citrated blood or disposable plastic, sterile "universal" container. The latter if AFB smear is required.	Do not refrigerate or pre- incubate inoculated bottles. Skin preparation is key to reducing contamination.

Authority For Issue: Benjamin Moore	Page 12 of 24
Document printed from Q-pulse 16/1	0/2023 09:28:00 by benjamin.moore

Document	SMRL65	Version No.	11	
No.		Issue Date	10/10/2023	

CSF	Disposable plastic, sterile "universal" container	Recommended minimum volume of 6mls. Smear and culture of 6mls CSF may have greater sensitivity than PCR.
Stools (not recommended)	Disposable plastic, sterile "universal" container	These are rarely informative
(minimum 1g)	with spoon	Not recommended.
PM specimens	Disposable plastic, sterile "universal" container or larger screw capped sterile container if	DO NOT USE FORMOL SALINE
		OR FORMALIN
Pus	Disposable plastic, sterile "universal" container Pus swab may be acceptable. Use 'Stuart's' transport media for pus swabs.	Pus is always preferred if available. Microscopy is not routinely performed on swabs.
Cultures	Egg medium slopes or aliquots of liquid medium	At least 5mls required if grown in liquid medium
Water and air from heater- cooler units (50 ml)	Sodium thiosulphate should added to the water collection pots to neutralise the hypochlorite before water sampling (final concentration of at least 18mg/L)	Air sampling plates supplied by SMRL by prior arrangement.

Authority For Issue: Benjamin Moore		Page 13 of 24
Document printed from Q-pulse 16/1	10/2023 09:28:00 by benjamin.moore	

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

#### Tests Available, Turn Around Times and Costs

Turnaround times for cultures are dependent on receipt of a pure culture containing sufficient mycobacteria for analysis.

The time interval between receipt of a culture and the issue of the final identification and first line sensitivity report varies greatly from 1-12 weeks, depending on factors such as the nature of the culture medium used by the sending laboratory, paucity of organisms in the culture, the species of Mycobacteria and the presence or absence of contamination.

Target turnaround times to result availability for 95% of samples:

Test	<b>Turnaround time in working days) (</b> 5d/ week laboratory service)
Primary sample AP smear microscopy	<1d from receipt of specimens (mean 0.3d)
Automated liquid and conventional solid culture on all samples being processed for mycobacterial culture	Set up ≤1 working day from receipt of specimens (Mean 1d)
Clinical specimen examination, culture, isolation and identification where the isolate is subsequently shown to be MTB complex	≤42d from receipt of specimens (mean 12.4d)
Provisional identification of MTB complex	≤7d from receipt of a positive cultures (mean 1.9d)
Identification of MTB complex by WGS	≤7d from receipt of a positive cultures (anticipated TAT subject to review)
Identification of NTM by WGS	≤7d from receipt of a positive cultures (anticipated TAT subject to review)

Authority For Issue: Benjamin Moore	Page 14 of 24
Document printed from Q-pulse 16/1	0/2023 09:28:00 by benjamin.moore

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

Sensitivity testing of MTB complex #	≤28d of isolate receipt/ isolation
	(mean 13d)
MTB complex relatedness by WGS	≤12d from receipt of isolate
	(anticipated TAT subject to review)
Rapid detection of MTB complex	≤3d from receipt of specimens
and detection of rifampicin resistance associated mutations on	(mean 1.9d)
clinical specimens	
Rapid detection of MTB complex	≤3d from receipt of specimens
and detection of isoniazid, fluoroquinolone, amikacin,	(anticipated TAT subject to review)
kanamycin, capreomycin and	
ethionamide resistance associated	
mutations	

# First line susceptibility results availability

The costs for NHS users are reviewed annually. These, plus costs for private tests are available on request.

Authority For Issue: Benjamin Moore	Page 15 of 24
Document printed from Q-pulse 16/1	0/2023 09:28:00 by benjamin.moore

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

Test	Method	Test frequency and turn	Comments
		around time	
Microscopy for acid-fast bacilli (AFB) detection	Auramine Phenol staining	Within 1 working day if receipted prior to mid-day (Mon-Fri)	All specimens examined by AP microscopy are also cultured for mycobacteria.
Primary Culture	Solid (modified LJ pyruvate egg slope) and rapid liquid culture (MGIT)	Daily. Usually available 2-8 weeks after receipt of sample. Cultures are reported negative at 8 weeks if appropriate but kept for 12 weeks in total.	Results from contaminated specimens may be delayed
Species identification from cultures	Whole Genome Sequencing.	Twice weekly (Mon/Thurs); 5-7 days	Rapid confirmation of MTB complex is achieved using lateral-flow immunochromatographic testing
Drug susceptibility testing of MTB comple	Whole Genome Sequencing.	Twice weekly (Mon/Thurs); 5-7 days	Some resistance predictions require confirmation by phenotypic testing which may delay some results. Phenotypic Tests for 3rd line TB drugs set up on request
Drug susceptibility testing of non- tuberculous myco "Slow Growers"	Broth microdilution	Set up weekly as required; 3-5 days incubation	For slow growing NTM mycobacteria. We are unable to test susceptibility to clofazamine.
Drug susceptibility testing of "Rapid Growers"	Broth microdilution	Set up weekly as required; 3-14 days incubation	For rapidly growing mycobacteria. We are unable to test susceptibility to clofazamine.
Rapid detection of MTB complex and detection of rifampicin resistance associated mutations	real time PCR (Cepheid)	Daily; usually available on day/day after receipt if received prior to mid-day.	Test is only validated for pulmonary samples although may provide useful for other specimen types. Pulmonary samples from NHS Lothian patients requesting TB investigation are routinely tested if they meet NICE based criteria: BAL samples (non ITU); HIV positive/significant immunosuppression; high index of TB suspicion; TB contact/ potential for large scale contact-screening; travel/residence in high incidence and/or MDR-TB country; on specific request by a consultant in respiratory/consultant infectious diseases; <15 years of age

Authority For Issue: Benjamin Moore	Page 16 of 24
Document printed from Q-pulse 16/	10/2023 09:28:00 by benjamin.moore

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

Rapid detection of MTB complex and detection of isoniazid, fluoroquinolone , amikacin, kanamycin, capreomycin and ethionamide resistance associated mutations	real time PCR (Cepheid)	Daily; usually available on day/day after receipt if received prior to mid-day.	Run by specific request of a consultant in respiratory/infectious diseases where there is a high index of suspicion for isoniazid mono-resistance, multidrug –resistant (MDR) or extensively drug- resistant (XDR) TB.
M. tuberculosis	WGS	Twice weekly (Mon/Thurs);	All new TB patients' isolates are sequenced.
typing (MTB		7-10 days	
Complex			
relatedness)			

#### **Molecular Detection of Mycobacteria in Clinical Specimens**

Note on rapid molecular tests: false negatives and false positives can and rarely do occur.

We currently use Xpert MTB/RIF Ultra (Cepheid) for the direct detection of MTBC in clinical specimens and for the rapid detection of rifampicin resistance associated mutations. This is performed on a daily basis for those NHS Lothian pulmonary patient specimens fulfilling NICE based criteria and for any AFB smear positives specimens.

For detection of isoniazid mono-resistance, MDR or XDR TB, we use Xpert MTB/XDR (Cepheid). This test can be run on both primary samples and positive cultures. Please note that this test detects a limited number of common mutations associated with drug resistance. Resistance will not be detected if alternative mutations occur. This test is tun by specific justification and request of a consultant in microbiology, respiratory or infectious diseases.

Authority For Issue: Benjamin Moore	Page 17 of 24
Document printed from Q-pulse 16/1	0/2023 09:28:00 by benjamin.moore

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

#### **Reporting of Results**

#### Policy and procedures for telephoned results

Previously unknown AFB smear positive or discrepant samples are telephoned as soon as possible, usually within 1 working day of receipt. New positive MTB complex cultures obtained from smear negative samples are also phoned as soon as possible after their presence is confirmed in the culture.

All NTM are reported and/or E-mailed. 'New' *M.abscessus* from NHS Lothian locations are telephoned in addition.

The following examples will also be telephoned or communicated urgently:

- results of direct rapid molecular tests on specimens if findings are unexpected.
- clinically important identification or drug susceptibility test results (e.g. primary resistance to first line drugs; acquired new resistance; unexpected mycobacterial species).
- new positive cultures from smear negative specimen/blood.

<u>Emailing of reports</u> to secure nhs.net accounts is set up already for all Scottish user laboratories. In these instances no written report is sent out. We can set this up if you have such a secure laboratory address to automatically email each time there is an updated report. Please contact Mike Smith or Mori Robertson in the first instance if you wish to use this means of reporting.

Authority For Issue: Benjamin Moore	Page 18 of 24
Document printed from Q-pulse 16/1	10/2023 09:28:00 by benjamin.moore

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

Written report formats

Initial Report on specimens				
Microscopy results				
Acid Fast Bacilli NOT seen				
Culture report to follow				
Microscopy results				
Acid Fast Bacilli seen (enumeration given)				
Culture report to follow				
Microscopy results				
Microscopy findings equivocal. Advise repeat s	sample.			
Equivocal result means 1-2 acid fast bacilli see	n per 300 fields. This is NOT a definite positive result			
and artefacts could cause this level of acid fast	t material in a specimen.			
Culture report to follow				
(approximately ½ such results are culture nega	ative)			
If rapid molecular testin	g is also performed on sample:			
<u>M. tuberculosis complex PCR</u>				
No MTB complex detected in this specimen or	n direct PCR.			
This does NOT rule out MTB complex or non-to	uberculous mycobacterial infection.			
<u>M. tuberculosis complex PCR</u>				
MTB complex DETECTED in this specimen on d	irect PCR, consistent with active infection but			
subject to culture and subsequent confirmator	ry results.			
No mutation indicating rifampicin resistance d	etected. NOTE:- This test is unable to detect true			
resistance to rifampicin in a small percentage	of isolates. This result is to be confirmed by isolation			
and susceptibility testing of the organism.				
<u>M. tuberculosis complex PCR</u>				
MTB complex DETECTED in this specimen on d	irect PCR, consistent with active infection but			
subject to culture and subsequent confirmatory results				
Mutations indicating resistance to rifampicin DETECTED				
This result is to be confirmed by isolation and susceptibility testing of the organism				
· ··· · · · · · · · · · · · · · · · ·				
<i>M. tuberculosis</i> complex PCR				
MTB complex DETECTED in this specimen on direct PCR consistent with active infection but				
subject to culture and subsequent confirmatory results				
Rifampicin resistance probe uninterpretable				
Authority For Issue: Benjamin Moore	Page 19 of 24			

Document printed from Q-pulse 16/10/2023 09:28:00 by benjamin.moore

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

#### M. tuberculosis complex PCR

M.tuberculosis complex probe uninterpretable.

#### **Initial Reports on Cultures received**

Confirmation as a Mycobacterium with species identification is normally reported within 7 working days of receipt of a culture. This is achieved by AFB microscopy to confirm the presence of mycobacteria and WGS for identification of cultures.

Initial rapid confirmation of MTB complex is achieved earlier using a lateral-flow immunochromatographic assay.

#### **Final Report on Cultures received**

Confirmed species identification with first line MTB complex sensitivity results may take from as little as 5 days to up to 6 weeks. This is achieved using WGS with confirmation of resistance predictions performed using phenotypic DST where appropriate.

Confirmed species identification of NTM may take 5-7 days. This is achieved using WGS. Phenotypic sensitivity testing if performed, is batched on a weekly basis.

*M. tuberculosis* complex genotyping results are reported when completed and comparisons of the genotypes of strains suspected to be clustered are reported.

#### **Cumulative reports**

Cumulative results on a patient with positive results can be produced on request. Please contact senior BMS or medical staff at SMRL.

At present only Authorised Final reports on each 'test' are available on SCI store. Interim results are not available.

Authority For Issue: Benjamin Moore		Page 20 of 24
Document printed from Q-pulse 16/7	10/2023 09:28:00 by benjamin.moore	

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

#### Technical, Scientific and Health and Safety Advice

The technical and scientific staff at the SMRL are happy to provide advice on all laboratory aspects of microscopy and isolation of Mycobacteria and on the safe running and design of containment level 3 laboratories.

Please contact Mr Mike Smith Mrs Pauline Claxton or Ms Ania Pilarska (BMS Team Managers') for technical help and advice.

#### **Clinical, Epidemiological and Infection Control Advice**

The primary source of medical advice for the diagnosis, management and public health investigation of TB lies with the consultants in the relevant specialties in the local health board. The consultant microbiologists attached to the SMRL are happy to complement this primary advice where appropriate.

Please contact Dr Ian Laurenson (Director), Dr Olga Moncayo and Dr Donald Inverarity (Deputy Directors) for advice on issues such as microbiological diagnostic tests, antibiotic susceptibility tests, strain typing for contact tracing, and infection control.

#### **Training and Education**

#### **Audit Study Day**

Every two years the laboratory hosts an Audit/Study day to allow SMRL staff to meet and discuss various issues with reference laboratory users, public health consultants, chest physicians and Health Protection Scotland (HPS) representatives.

#### Attachments and visits

Each year SMRL provides training attachments for staff from other laboratories including Biomedical Scientists, Clinical Scientists and Trainee Doctors. SMRL also provides some support for undergraduate science and medical students, MSc and PhD students and projects for Trainee Doctors.

#### Visits and lectures by SMRL staff to other institutions

Staff at SMRL provide teaching and advice at local and national level on invitation particularly in the areas of diagnosis, patient management and molecular epidemiology.

#### **Contact details**

To discuss short visits to SMRL for training purposes, please contact Mr. Mike Smith, BMS Team Manager or Mrs Mori Robertson, Reference Laboratory Administrator in the first instance.

Authority For Issue: Benjamin Moore	Page 21 of 24
Document printed from Q-pulse 16/1	0/2023 09:28:00 by benjamin.moore

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

#### **Research and Audit**

Staff are willing to collaborate in relevant audit and research activities. This may take the form of simple data searches to more formal grant applications dependent on ethical approval where relevant and availability of resources.

#### **Contact details**

Please contact Dr Ian Laurenson (Director), Dr Olga Moncayo (Deputy Director) or Dr Louise Seagar (Clinical Scientist) for discussions regarding research/audit collaboration.

#### **Quality Assurance**

The SMRL is committed to high quality laboratory analysis and is accredited by the United Kingdom Accreditation Service to ISO 15189:2012 for the scope of testing. Environmental water sampling, susceptibility testing to non-tuberculosis mycobacteria, direct molecular detection of non pulmonary samples is currently out with our scope of accreditation.

The laboratory participates in the UK National External Quality Assurance (NEQAS) scheme for Mycobacteria microscopy, culture and molecular typing as well as Quality Control for Molecular Diagnostics (QCMD) for molecular detection in specimens. The laboratory also participates in the WHO 'Drug Proficiency Testing Scheme', an external QA scheme for sensitivity testing of *M tuberculosis* isolates. The laboratory also participates in the INSTAND Mycobacteria Drug Susceptibility Testing of Non-tuberculous Mycobacteria EQA program. For typing, the laboratory partakes in the international proficiency study on WGS by the National Institute for Public Health and the Environment TB Reference Laboratory Netherlands (RIVM).

We also carry out Internal Quality Assurance (IQA).

The SMRL Annual Report and User Manual are available on Scottish Microbiology Reference Laboratories, Edinburgh. <u>http://www.edinburghlabmed.co.uk/Specialities/reflab/Pages/default.aspx</u>

(accessed January 2022)

#### **Protection of Personal Information**

The SMRL and NHS Lothian Mycobacteria Laboratory comply with the Data Protection Act. This requires all organisations which handle personal information to comply with a number of important principles regarding privacy and disclosure.

Authority For Issue: Benjamin Moore	Page 22 of 24
Document printed from Q-pulse 16/1	0/2023 09:28:00 by benjamin.moore

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

#### **Antimicrobial Assays**

The following information is for assays on first line TB therapy drugs. Information on other anti mycobacterial drugs can be found at <u>http://www.assayfinder.com/</u> (accessed January2022)

·····	
Provider	Cardiff Toxicology Laboratories
Contact Name	Katie Jones, Clinical Scientist, katie.jones14@wales.nhs.uk
Address For Specimens	Toxicology Laboratory The Academic Centre Llandough Hospital Penarth CF64 2XX
Country	United Kingdom
Tel. No. (Voice)	029 2071 6894
Tel. No. (Fax)	029 2035 0142
E Mail (click to mail)	Katie.Jones14@wales.nhs.uk
Accreditation Status	Accredited
Other Information	For general enquiries please contact Emma.Taylor6@wales.nhs.uk
Web Site (click to view)	http://www.ctlabs.co.uk

Isoniazid, Pyrazinamide and Ethambutol

#### **Rifampicin and Moxifloxacin**

Provider	Antimicrobial Reference Laboratory, Southmead Hospital, Bristol
Contact Name	Alan Noel
Address For Specimens	Department of Medical Microbiology Southmead Hospital Westbury on Trym Bristol BS10 5NB
Country	United Kingdom
Tel. No. (Voice)	0117 4146220
Tel. No. (Fax)	0117 4146282
E Mail (click to mail)	arlenquiries@nbt.nhs.uk
Accreditation Status	Accredited
Other Information	
Web Site (click to view)	Requesting   North Bristol NHS Trust (nbt.nhs.uk)

Authority For Issue: Benjamin Moore	Page 23 of 24
Document printed from Q-pulse 16/1	0/2023 09:28:00 by benjamin.moore

Document	SMRL65	Version No.	11
No.		Issue Date	10/10/2023

#### **Leprosy Diagnosis**

The SMRL does not undertake investigations for the diagnosis for leprosy.

Clinicians should refer all suspected or confirmed cases to one of the national leprosy centres where the case can be reviewed by a Consultant Advisor in Leprosy. Contact in the first instance:

Dr. Stephen Walker Consultant Advisor in Leprosy Hospital for Tropical Diseases Mortimer Market Centre Capper Street London WC1E 6JB Tel: 020 3447 5959

Detection of *M. leprae* by PCR is available at Great Ormond Street Hospital. Samples should be sent directly by the user to:

Microbiology, Virology and Infection Control Level 4, Camelia Botnar Laboratories Great Ormond Street Hospital for Children Great Ormond Street London WC1N 3JH DX number: DX6640203 DX Exchange: Bloomsbury 91WC

Authority For Issue: Benjamin Moore		Page 24 of 24
Document printed from Q-pulse 16/7	10/2023 09:28:00 by benjamin.moore	